

RESEARCH ARTICLE



A Study to Determine the Effect of Egg Albumin Dressing On Peristomal Wound Healing of the Colostomy Patients in a Selected Hospital, Kolkata, West Bengal

Ms. Rimi Chakraborty¹ | Dr Arpan Dutta Roy² | Dr Sayantan Ghosh^{3*} | Dr Pankaj Kumar Singh⁴ | Dr. Prolay Paul⁵

¹M Sc. Critical Care Nursing
WBUHS, Emergency and Disaster
Nursing (INC, New Delhi), BLS,
ACLS& PALS Instructor,
American Heart and Stroke
Association MBA Hospital
Administration, ICFAI, FCCS,
Society of Critical Care Medicine
(USA) NABH Internal Assessor
(POI of NABH 5th Edition), QCI,
CPHIC in Basic Infection Control,
CAHO

²Chief of Clinical Pharmacology
Ruby General Hospital Certified
NABH Assessor (QCI) HOPE
Doctor of Pharmacy (PB) Clinical
Research, (PhD.) Pharmacology
Certified GCP, ACLS,
Pharmacovigilance, NABH
assessor

³Pharm D Intern Department of
Pharmacy Practice Sri
Adichunchanagiri College of
Pharmacy, Karnataka

⁴MBBS, DNB Emergency Medicine
Attending Consultant Department
of Emergency Medicine Ruby
General Hospital Kolkata

⁵Pharm D Intern Department of
Pharmacy Practice Sri
Adichunchanagiri College of
Pharmacy Karnataka

Abstract

The researcher conducted a quasi experimental study to evaluate the effect of egg albumin dressing on peristomal wound healing in a selected hospital, Kolkata, with the objectives to assess the peristomal skin condition of colostomy patients before treatment, to evaluate the effect of egg albumin dressing on healing of peristomal area and reduction of pain, to find out the association between the peristomal wound healing and selected variables. The final study was conducted at Curzon ward, Victoria ward of SSKM hospital, Kolkata. Ethical permission was sought out from Ethical Committee of SSKM hospital, Kolkata. Informed consent was taken from all respondents. The sample was selected according to their selected criteria. The sample selection was done by purposive sampling. They were randomly assigned into two groups (experimental and control group) in 1:1 ratio. The study concluded with its limitations, implications and recommendations for conducting a study may be conducted for a longer duration of observation with the treatment.

Keywords: Albumin Dressing, Peristomal Wound Healing, Colostomy Patients, Egg Albumin

Copyright : © 2021 The Authors. Published by Publisher. This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).



1 | INTRODUCTION

According to the WHO reports in 2012, cancer is a leading cause of death worldwide, accounting for 8.2 million deaths. Amongst the most common causes of cancer death, about 694000 are from colorectal cancer. Most of the operable colorectal cancers require a surgical procedure called colostomy.

A colostomy is major surgery that creates an opening (known as a “stoma”) in the colon to permit waste to exit outside the body into a pouch attached to the abdomen. Generally, in a colostomy, part of or the entire colon is removed. A colostomy may be permanent or temporary, depending on the medical condition that has necessitated the surgery. A permanent colostomy is customarily performed when the rectum or most of the colon is removed. (1)

The main purposes of a wound dressing are, to clean the site, absorb exudates, if any, ease pain and provide protection from infection. The wound dressing should ideally fulfill some primary and secondary requirements. (2)

The primary requirements would be that the dressing is free of toxic or irritant extractable, should not release particles or non-biodegradable fibres into the wound, should form an effective bacterial barrier, forms an effective water-resistant seal to the periwound skin, but is easily removable without causing trauma or skin stripping should be able to maintain the wound and the surrounding skin in an optimum state of hydration, provide protection to the periwound skin from potentially irritant wound exudates and excess moisture, produce minimal pain during application or removal as a result of adherence to the wound surface and maintain the wound at the optimum temperature and pH. The secondary requirements should include antimicrobial activity, ability to remove or inactivate proteolytic enzymes in chronic wound fluid, possess haemostatic activity and have effective wound debriding activity. (3)

2 | NEED OF THE STUDY

One of the main types of stoma is colostomy, which has a risk of forming sore on the peristomal skin. Through the stomas, feces and body fluid are collected in the stoma appliance. The stoma appliance is attached to the peristomal skin with adhesive. As there is a chance of continuous seepage of feces and body fluid through this stoma there is a high chance of skin excoriation at peristomal region due to the corrosiveness of that feces and body fluid. Also the continuous pressure and friction caused due to the adhesive of the stoma appliance contribute to the chances of excoriation of the peristomal skin. The severity of the excoriation depends primarily on these factors. It is essential to ensure that the skin surface, on which the appliance is attached, is free from breaks or soreness as this might lead to appliance leakage.

Generally enterostomal therapist takes care of these stomas in the post operative period. As there is an inadequate number of enterostomal therapist, the general nurse also has a vital role to take care of the patient with a stoma. Taking care of the patient with any ostomy is indeed a challenge to any nurse.

The investigator during her clinical experience noticed that the peristomal skin excoriation is very common in the patient having colostomy and different types of dressing, commercially available in the market viz, ostomy powder, ostomy paste, hydrocolloid based appliances, etc are applied to reduce peristomal skin complications. The investigator, considering the increasing number of cases from different economic backgrounds, has felt that there is a need to look into alternative dressing materials.

Use of egg white for treatment and healing of wounds was an old Roman technique for treating gunshot wounds. Egg white constitutes about 20-25% of the

Supplementary information The online version of this article (<https://doi.org/10.15520/jcmro.v4i04.409>) contains supplementary material, which is available to authorized users.

Corresponding Author: *Dr Sayantan Ghosh*
Pharm D Intern Department of Pharmacy Practice Sri
Adichunchanagiri College of Pharmacy, Karnataka
Email: gsayantan26@gmail.com

egg. The egg white is composed of proteins and minerals. Different types of proteins are present in egg white.

Some of them are Ovalbumin, Conalbumin, Ovumucoid, Ovomucin, Lysozyme, Avidin,

Ovoglobulin, Ovoinhibitor. It also contains minerals like Sulphur, which has antibacterial and anti-inflammatory properties and Copper which is toxic to bacteria and also used in a number of rejuvenating and skin revitalizing treatments. These properties of egg albumin make it suitable to be used in topical application in medical dressing.

Thus the investigator thought that topical application of egg white dressing may be an effective healing agent for peristomal wounds.

3 | OBJECTIVES –

- To evaluate the effect of egg albumin dressing on healing of the peristomal skin area and reduction of pain among experimental group of colostomy patients.
- To assess the peristomal skin condition among experimental group of colostomy patients before treatment.
- To assess the peristomal skin condition among control group of colostomy patients before treatment.
- To find out the association between the peristomal wound healing and selected sample characteristics.

4 | STUDY CRITERIA-

Inclusion Criteria-

- Colostomy patients admitted in the surgical ward on their 5th postoperative day onward
- Patients who are willing to participate in the study

- Adult patient >18years of age irrespective of their disease condition.

Exclusion Criteria-

- Known allergic condition to egg albumin

5 | MATERIALS & METHOD-

Study Type-

The Study was a Quasi experimental research approach.

Study Design

The design adopted for this study is pre-test post-test control group time series design.

Operational Definitions-

Colostomy Patient - In this study, colostomy patient refers to patients more than 18 years age, admitted in the surgical ward of the selected hospital on the 4th postoperative day of permanent or temporary colostomy.

Peristomal skin - It refers to the area surrounding the stoma where appliance is attached.

Egg albumin dressing - It refers to the direct application of the raw egg white portion with the help of sterile gauge piece, once in a day, on alternate days, for a total of three times, on the peristomal skin, after cleaning the region with 0.9% normal saline

Effect – It refers to whether the desired effect of egg albumin dressing has achieved or not and is measured by healing score.

Peristomal skin wound - Peristomal skin wound is assessed by modified Ostomy Skin Tool, the wound status is assessed through rating scale and will be measured by DET Scoring in terms of the discoloration, erosion, and tissue overgrowth. Table 1

Selected sample characteristics - Selected sample characteristics will include demographic profile (consist of age, sex, education, occupation, income), health assessment (height, weight BMI), illness profile (duration of illness, time taken to diagnose, duration of peristomal skin wound, cancer stage, nature of surgery, no of postoperative days in intensive care

TABLE 1: Schematic representation of data collection instruments -

Tool No	Name of the tool	Variables to be measured	Technique
Tool-I	Semi-structured interview schedule	Demographic profile	Interview
Tool-II	Health assessment proforma analysis proforma	Height, weight, BMI Illness profile	Measurement Record analysis
Tool-III	Modified Ostomy Skin Tool	Peristomal skin wound status	Assessment
Toll-IV	Visual analogue scale	Wound pain	Assessment

unit, presence of Diabetes mellitus, hypertension, feeding pattern, blood report of HB%, WBC, ESR).

Data Collection Procedure-

The final study was conducted at Curzon ward, Victoria ward of SSKM hospital. Kolkata.

Ethical permission was sought out from Ethical Committee of SSKM hospital, Kolkata. Informed consent was taken from all respondents. The sample was selected according to their selected criteria. The sample selection was done by purposive sampling. But randomly assigned into two groups (experimental and control group) in 1:1 ratio. First one was selected as experimental group and second one as control group. In this way 15 patients in the experimental group were selected and coded as E1, E2, E3, E15 and another 15 patients in the control group were selected and coded as C1, C2, C3 C15.

The Data was Analysed-using

Section I- The findings related to the description of the demographic characteristics of the colostomy patients presented in frequencies and percentage distribution.

Section II A- The findings related to the description of the health assessment of the colostomy patients presented in frequencies and percentage distribution.

Section II B- The findings related to the description of the illness profile of the colostomy patients presented in frequencies and percentage distribution.

Section III- The findings related to the pre intervention score of experimental group and control group by mean, median, and standard deviation.

Study Site

The study was conducted at the surgical ward of the SSKM Hospital, Kolkata.

6 | RESULTS-

Tables 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 and 18

Section-II A Finding related to the description of the health assessment of the colostomy patients

Section-III Findings related to the pre intervention peristomal skin wound score of experimental group and control group by mean, mean difference and standard deviation.

Section-IV Finding related to the distribution of the colostomy patients according to the preobservation pain score observed by VAS

Section V Effectiveness of egg albumin dressing for Peristomal skin wound healing.

There is a significant difference of mean score of peristomal skin wound discoloration in colostomy patients in experimental group before and after application of egg albumin dressing as measured by modified ostomy skin tool at 0.05 level of significance.

There is a significant difference of mean score of peristomal skin wound erosion in colostomy patients in experimental group before and after application of egg albumin dressing as measured by modified Ostomy Skin Tool at 0.05 level of significance.

There is a significant difference of mean score of peristomal skin wound tissue overgrowth in colostomy patients in experimental group before and

TABLE 2: Frequency and percentage distribution of the colostomy patients according to their age, sex and educational qualification. n = 30(15+15)

Sample Characteristics	Experimental Group		Control Group	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
Age (In years)				
18-30	1	6.7	1	6.7
31-50	10	66.7	11	73.3
51-70	4	26.7	3	20
Sex				
Male	8	53.3	10	66.7
Female	7	46.7	5	33.3
Educational Qualification				
Primary	8	53.3	9	60.0
Secondary	5	33.3	3	20.0
Higher Secondary & Above	2	13.3	3	20.0

TABLE 3: Frequency and percentage distribution of the colostomy patients according to their occupation, monthly family income and diagnosis n = 30(15+15)

Sample Characteristics	Experimental Group		Control Group	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
Occupation				
Service	2	13.3	3	20
Daily labor	5	33.3	4	26.7
Business	8	53.3	8	53.3
Monthly Family Income				
< 4000/-	Nil	-	1	6.7
4000/- and above	15	100	14	93.3
Diagnosis				
CA Rectum	7	46.7	6	40
CA Colon	8	53.3	9	60

after application of egg albumin dressing as measured by modified Ostomy Skin Tool at 0.05 level of significance.

There is a significant difference of mean post intervention score of Peristomal skin wound discoloration in colostomy patients in the experimental group getting egg albumin dressing than that of control group assumed to get conventional treatment at 0.05 level of significance.

There is a significant difference of mean post intervention score of peristomal skin wound erosion among colostomy patients in the experimental group

getting egg albumin dressing than that of control group assumed to get conventional treatment at 0.05 level of significance.

There is a significant difference of mean post intervention score of Peristomal skin wound tissue overgrowth among colostomy patients in the experimental group getting egg albumin dressing than that of control group assumed to get conventional treatment at 0.05 level of significance.

Section VI: Effectiveness of egg albumin dressing for reducing degree of Peristomal skin wound pain.

TABLE 4: Frequency and percentage distribution of colostomy patients according to their height, weight and BMI

Sample Characteristics	Experimental Group		Control Group	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
Height in cms				
>137-150	6	40	7	46.7
>150-163	7	46.7	6	40
>163-176	2	13.3	2	13.3
Weight in kg				
40-45	4	26.7	5	33.3
45-50	7	46.7	5	33.3
50-55	3	20	3	20
55-60	Nil	-	1	6.6
60-65	1	6.6	1	6.6
BMI				
Normal	10	66.7	9	60
Low	5	33.3	6	40

TABLE 5: Frequency and percentage distribution of colostomy patients according to their cancer stage, type of surgery performed and no of days stay in ICU

Sample Characteristics	Experimental Group		Control Group	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
Cancer stage				
Stage-I	5	33.3	6	40
Stage-II	6	40	6	40
Stage-III	4	26.7	3	20
Types of surgery performed				
Therapeutic	15	100	15	100
Palliative	Nil	-	Nil	-
No of days stay in ICU				
2 Day	Nil	-	2	13.3
3 Day	Nil	-	Nil	-
None	15	100	13	86.7

There is a significant difference of mean Peristomal skin wound pain score among colostomy patients in experimental group before and after application of egg albumin dressing as measured by VAS at 0.05 levels of significance.

There is a significant difference of mean Peristomal skin wound pain score among colostomy patients in experimental group before and after application of

egg albumin dressing as measured by VAS at 0.05 levels of significance.

There is a significant difference of mean post intervention score of pain among colostomy patients in the experimental group getting egg albumin dressing than that of control group assumed to get conventional treatment at 0.05 levels of significance.

TABLE 6: Frequency and percentage distribution of colostomy patients according to the presence of diabetes mellitus, hypertension and haemoglobin levels

Sample Characteristics	Experimental Group		Control Group	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
Diabetes mellitus				
Absent	11	73.3	9	60
Present	4	26.6	6	40
Hypertension				
Absent	15	100	12	80
Present	Nil	-	3	20
Hb level				
Normal	7	46.7	6	40
Below normal	8	53.3	9	60

TABLE 7: Frequency and percentage distribution of colostomy patients according to their WBC count, ESR level and mode of feeding

Sample characteristics	Experimental Group		Control Group	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
White blood cell count				
Above normal	4	26.7	7	46.7
Normal	11	73.3	8	53.3
ESR level				
Above normal	Nil	-	Nil	-
Normal	15	100	15	100
Mode of feeding				
Enteral	15	100	15	100
Parental	Nil	-	Nil	-

TABLE 8: Mean, Mean Difference and Standard Deviation of pre-intervention score of experimental and control group of colostomy patients

Group	Domain	Mean	Mean D	SD
Experimental	Discolorati	4.06	0	0.99
Control		4.06		0.99
Experimental	Erosion	3.80	0.07	0.97
Control		3.73		0.98
Experimental	Tissue over-growth	2.26	0.07	0.92
Control		2.33		0.93

TABLE 9: Frequency and percentage distribution of colostomy patients according to the preintervention pain score observed by VAS

Degree of Pain	Experimental Group		Control Group	
	Frequency (f)	Percentage (%)	Frequency (f)	Percentage (%)
Little Discomfort (1-2)	1	6.7	1	6.7
Mild Pain (3-4)	5	33.3	5	33.3
Moderate Pain (5-6)	8	53.3	7	46.7
Severe pain (7-8)	1	6.7	2	13.3

TABLE 10: Mean, Mean Difference, SD, SE and Paired 't' value of pre and post intervention colostomy skin wound discoloration score by modified Ostomy Skin Tool in experimental Group n = 15

Observation	Mean	MD	SD	SE	't'
Before treatment	4.06	2.06	0.99	0.42	4.84*
After treatment	2.00		1.31		

df (14) = 2.15, p < 0.05

TABLE 11: Mean, Mean Difference, SD, SE and Paired 't' value of pre and post intervention colostomy skin wound erosion score by modified Ostomy Skin Tool in experimental n = 15

Observation	Mean	MD	SD	SE	't'
Before treatment	3.8	2.00	0.97	0.42	4.69*
After treatment	1.8		1.32		

df (14) = 2.15, P < 0.05

TABLE 12: Mean, Mean Difference, SD, SE and Paired 't' value of pre and post intervention colostomy skin wound tissue overgrowth score by modified Ostomy Skin Tool in experimental Group

Observation	Mean	MD	SD	SE	't'
Before treatment	2.26	1.66	0.92	0.28	5.80*
After treatment	0.6		0.61		

df (14) = 2.15, p < 0.05

TABLE 13: Mean, Mean Difference, SD, SE and Unpaired 't' value of experimental and control group post observation score of discoloration measured by Modified Ostomy Skin Tool

Group	Mean	MD	SD	SE	't'
Experimental Group	2.0	-2.0	1.31	0.42	4.74*
Control Group	4.0		0.96		

df (28) = 2.05, p < 0.05

TABLE 14: Mean, Mean Difference, SD and Unpaired 't' value of experimental and control group post observation score of erosion measured by modified Ostomy Skin n=30(15+15)

Variable	Mean	MD	SD	SE	't'
Experimental Group	1.8	-1.8	1.32	0.43	4.16*
Control Group	3.6		1.01		

df (28) = 2.05, p < 0.05

TABLE 15: Mean, Mean Difference, SD and Unpaired ``t'' value of experimental and control group post observation score of tissue overgrowth measured by modified Ostomy Skin n = 30 (15+15)

Variable	Mean	MD	SD	SE	``t''
Experimental Group	0.6	-1.4	0.61	0.26	5.31*
Control Group	2.00		0.81		

df (28) = 2.05, p < 0.05

TABLE 16: Mean, Mean Difference, SD and Paired ``t'' value of pre and post Intervention peristomal skin wound pain score by VAS in experimental Group n = 15

Observation	Mean	MD	SD	SE	``t''
Before treatment	5.2	4.0	1.42	0.48	8.25*
After treatment	1.2		1.22		

df (14) = 2.15, P < 0.05

There is a significant difference of mean post intervention score of pain among colostomy patients in the experimental group getting egg albumin dressing than that of control group assumed to get conventional treatment at 0.05 levels of significance..

Section-VII Findings related to the association between peristomal skin wound healing and illness profile of the colostomy patients.

7 | DISCUSSION AND CONCLUSION:

The present study was likely to be supported by the study of Parkinson, 1999, who conducted a study and evaluated that the major proteins of albumen are ovalbumin, conalbumin (ovotransferrin), ovomucoid, lysozyme and ovomucin. Lysozyme which forms a chemical protection against microorganism, by dissolving the cell wall of bacteria, constitutes about 3.5% of the egg. This prompted the researcher to conduct this study with a desire to study the effect of application of egg albumin, in peristomal wound dressing, with respect to its healing of the wound.

The present study has revealed that satisfactory healing of the peristomal skin wound was achieved by the application of egg albumin dressing in terms of reduction of irritation in the wound area. Similar reports have been published in the study to determine the effect of cyanoacrylate protectant to manage peristomal skin irritation under ostomy skin barrier wafers conducted by Catherine T. Milne, Darlene

Saucier, Chenel Trevellini, Juliet Smith (2010). Additionally, their study also reported the adhesive properties of egg albumin which helped in effective sealing between the stoma appliance and the peristomal skin.

The present study has also revealed that healing of the peristomal skin wound was achieved by the application of egg albumin dressing in terms of tissue overgrowth, discolouration, and controlling tissue erosion. This is line with the reports published by Zou, C, Kobayshi, K and Kato (1991) who had observed the morphological changes in some cell types under the influence of egg white, suggesting that egg white may promote cell differentiation.

REFERENCES

1. Salvadalena GD. The incidence of stoma and peristomal complication during the first 3 months after stoma creation. *Journal of Wound Ostomy Continence Nurse* 2013 Jul-Aug; 40 (4): 400-6 ;.
2. Adham M, Abdou, Mujo K, Kenji S. Functional Proteins and Peptides of Hen's Egg Origin. *Bioactive Food Peptides in Health and Disease* 2013; ISBN: 978-953-51-0964-8. ;.
3. Robert E, Feeney and David A. The antibacterial activity of the egg white protein conalbumin. *Journal of Bacteriology* 1952 Nov; 64(5): 629-43. ;.

TABLE 17: Mean, Mean Difference, SD, SE and Unpaired 't' value of experimental and control group wound pain score measured by VAS n=30

Group	Mean	MD	SD	SE	't'
Experimental	1.2		1.22	0.45	6.16*
Control	4.0	-2.8	1.26		

df (28) = 2.05, P < 0.05

TABLE 18: Chi-square to find out the association between selected sample characteristics and Peristomal skin wound healing among colostomy patients n =15

Sample characteristics	Post observation score		x2 value
	< Median	≥ Median	
BMI	8	2	2.8
Normal	1	4	
Below normal			
Diabetic Mellitus	2	2	1.03
Present	1	10	
Absent			
Haemoglobin Level	3	5	1.87
Below normal	6	1	
Normal			
WBC Count	1 10	3	1.83
Above normal		1	
Normal			

How to cite this article: Chakraborty M.R., D.A.D.R., Ghosh D.S., Singh D.P.K., Paul D.P. A Study to Determine the Effect of Egg Albumin Dressing On Peristomal Wound Healing of the Colostomy Patients in a Selected Hospital, Kolkata, West Bengal. Journal of Current Medical Research and Opinion. 2021;886–894. <https://doi.org/10.15520/jcmro.v4i04.409>