



Original Research

Nurses' and Midwives' Perceived Barriers to Implementing Kangaroo Mother Care in Kirkuk City Hospitals

Hind Ali Raji¹, Dr. Rabab Hamody Hanon²

¹Salah al-Din Health
Directorate / Tuz Health Sector

²College of Nursing University of
Kirkuk

Abstract:

Background: Kangaroo mother care is an evidence-based practice for preterm and low-birth-weight newborns that has been shown to reduce neonatal mortality and morbidity. Even though kangaroo mother care is beneficial, it is not widely implemented. Routine practices often continue despite barriers at the institutional, healthcare worker, maternal, and family levels.

Aim: To identify nurses' and midwives' perceived barriers to implementing Kangaroo Mother Care in Kirkuk City Hospitals.

Methods: A descriptive cross-sectional study was conducted in hospitals in Kirkuk City. 163 nurses and midwives were selected using a purposive non-probability sampling strategy. A standardized, self-administered questionnaire was used to gather data. Descriptive statistical analyses were performed on the data using SPSS version 26.0.

Results: A large percentage of nurses and midwives reported a high level of perceived barriers to KMC implementation (70.6%). Lack of professional experience or knowledge was the most frequently expressed barrier (86.5%). There were also reports of system-related barriers, including high workload (80.4%) and inadequate support (81%).

Conclusion: Implementing kangaroo mother care presents significant challenges for nurses and midwives, with 70.6% reporting a "high" level of perceived barriers.

Keywords: Nurses and Midwives, Perceived Barriers, Kangaroo Mother Care



Corresponding Author: **Hind Ali Raji**, Salah al-Din Health Directorate / Tuz Health Sector

Email Id: nsm24014@uokirkuk.edu.iq



©Copyright 2026 The Current Medical Research and Opinion. Licenced by Creative Commons Attribution-Non Commercial-No Derivatives (CC BY-NC-ND) 4.0 International License. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Received: 10.06.2026 | **Revised:** 14.07.2026 | **Accepted:** 20.07.2026 | **Published:** 01-08.2026

Introduction

Low birth weight and preterm birth (LBW and PTB) remain serious global public health problems that greatly increase the prevalence of disease (Liu et al., 2024). Approximately 15% of births worldwide are preterm, and preterm birth is a major cause of neonatal morbidity and mortality (Darghiyan et al., 2023). However, recent statistics indicate that Iraq's neonatal mortality rate improved from 14 deaths per 1,000 live births in 2021 to 13 deaths per 1,000 live births in 2022. In comparison, the global average stands at 12 deaths per 1,000 live births. These figures underscore the urgent need for advancements in newborn healthcare. (UNICEF, 2022). Kangaroo Mother Care (KMC) is a method that has received clinical validation and demonstrated efficacy in improving the health and survival of these vulnerable infants. Additionally, infants who received KMC—which involves direct contact between mother and infant—showed better weight gain and strengthened the mother-infant bond (Chamhene & Moshi, 2023). Kangaroo Mother Care also enhances maternal health in high-income, middle-income, and low-income countries. Compared to standard care, it significantly decreases hospital stays, hypothermia, hypoglycemia, sepsis, and neonatal mortality (Gebeyehu et al., 2022). Kangaroo Mother Care (KMC) was a program created in Colombia to help families care for their premature and low-birth-weight newborns (Abadía-Barrero, 2018). Although these principles are broadly acknowledged in clinical settings,

Methodology

Study design and setting

A descriptive cross-sectional study was conducted. The study was carried out on the midwives and nurses who work in the maternity units (delivery rooms) and the neonatal intensive care units of the three hospitals under the Kirkuk Health Directorate: Kirkuk Teaching Hospital, Azadi Teaching Hospital, and AL-Naser Maternity and Child Hospital.

their implementation within practical policies faces professional and administrative challenges (Maniago et al., 2020). To ensure that infants and their families receive thorough, patient-centered, and compassionate care, effective nursing practices are essential (Sales et al., 2018). Even though kangaroo mother care has demonstrated benefits, it is not commonly used. Routine procedures frequently make it difficult to apply it consistently, with barriers at the institutional, healthcare worker, maternal, and family levels. The majority of these barriers have more to do with established routines than with health issues (Alenchery et al., 2018). Institutional barriers include a lack of beds, space, and staff trained to provide clinical care, counseling, and family support. However, only a limited number of countries have successfully expanded the application of kangaroo mother care beyond the designated areas. (Calibo et al., 2021). According to previous studies, medical professionals and caregivers often report several reasons for their resistance to participating in KMC, including time constraints, cultural barriers, a lack of support from nursing staff, and a lack of understanding of the technique (Mala & Abdul-Sahib, 2022).

However, despite their potential significance in this understudied research environment, barriers to the application of Kangaroo Mother Care (KMC) in Kirkuk City have not yet been fully investigated. Due to regional variations in healthcare systems and cultural influences, there is a significant knowledge gap in this context.

Sample

A purposive non-probability sampling technique was used to select 163 nurses and midwives from the neonatal intensive care units and the maternity units (delivery rooms).

Inclusion Criteria

A. Nurses and midwives who provided direct care in the delivery room.

B. Nurses and midwives who provided direct care in the Neonatal Intensive Care Unit (NICU).

C. Nurses and midwives who covered all educational levels.

D. Nurses and midwives who had at least six months of experience in the delivery room and the Neonatal Intensive Care Unit (NICU).

Exclusion Criteria

A. Nurses and midwives who held administrative positions only, such as head nurses.

B. Nurses and midwives who were not present or on leave during the data collection period.

C. Nurses and midwives who refused to participate in the study.

Method of Data Collection

A structured questionnaire was used to collect the data.

The questionnaire was developed based on a comprehensive review of the literature and previous studies relevant to the study topic (Al-Shehri & Binmanee, 2021; Chan et al., 2017).

The questionnaire consisted of four parts:

Part (1): Sociodemographic Characteristics of Nurses and Midwives.

It consisted of five questions about the research participants' sociodemographic characteristics, including their age, marital status, level of education, place of residence, and socioeconomic status.

Part (2): Professional Characteristics:

It consisted of five questions that addressed the professional characteristics of midwives and nurses. These included their length of service, the number of years they worked in the intensive care

unit and delivery room, participation in training related to Kangaroo Mother Care, their work area, and their work shift.

Part (3): Knowledge of Nurses and Midwives Regarding Kangaroo Mother Care: It consisted of sixteen questions that covered physiological regulation in neonates, breastfeeding, immunity, and the clinical benefits of KMC.

Part (4): Barriers to the Implementation of Kangaroo Mother Care: It consisted of seventeen questions that included institutional and healthcare worker-related barriers and environmental or clinical barriers.

Validity and Reliability of the Study Instrument

A panel of ten experts evaluated the study questionnaire's content validity. The panel included three faculty members from the College of Nursing, University of Kirkuk; two from the College of Nursing, University of Mosul; two from the University of Baghdad; two pediatricians from hospitals in Kirkuk and Tikrit; and one faculty member from the College of Health Sciences, Hawler Medical University. Internal consistency was evaluated using Cronbach's alpha.

Data Analysis

The Statistical Package for the Social Sciences (SPSS), version 26.0, was used to analyze and interpret the data. Descriptive statistics included the mean (M), standard deviation (SD), frequency (f), and percentage (%).

Result

Table 1: Description of Participants' Sociodemographic Variables (SDVs) (N=163)

List	SDVs	No	%	
1	Age (Years) M±SD= 30 ± 9.8	20 – 29	94	57.7
		30 – 39	51	31.3
		40 – 49	12	7.3
		50 +	6	3.7
2	Marital status	Single	52	31.9
		Married	106	65
		Divorced	3	1.8
		Widowed	2	1.2
3	Level of education in nursing	Secondary school	10	6.1
		Midwifery Secondary sch.	22	13.4
		High Health Institution	93	57.1
		Diploma	26	16
		Bachelor	12	7.4
4	Residency	Urban	156	95.7
		Rural	7	4.3
5	Perceived Socioeconomic status	Sufficient	86	52.8
		Barely sufficient	59	36.2
		Insufficient	18	11

No.: Number, %: Percentage, M: Mean, SD: Standard deviation

Table 1 describes sociodemographic variables of nurses and midwives who participated in the study (N=163); the findings indicate that the average age is 30±9.8 years, and more than half (57.7%) of them fell within the age group of 20 – 29 years. The marital status reveals that a higher proportion (65%) were married. Regarding the level of education in nursing, more than half (57.4%) of

nurses and midwives graduated from high-quality institutions. The residency indicates that the majority (95.7%) of nurses and midwives were residing in urban areas. Concerning socioeconomic status, 52.8% of nurses and midwives perceive their socioeconomic status as sufficient.

Table 2: Description of Participants' Professional Variables (N = 163)

List	Variables	No	%
1	Years of experience M±SD= 7 ± 6	1 – 5	49.1
		6 – 10	30.7
		11 – 15	14.7
		16 – 20	1.2
		21 +	4.3
2	Current unit years of experience M±SD= 5 ± 5.5	> 1	3.7
		1 – 5	61.3
		6 – 10	20.2
		11 – 15	11
		16 – 20	1.2
		21 +	2.5
3	Participation in training courses	None	55.2
		1 – 3	33.7
		4 – 6	8.6
		7 – 10	2.5
4	Current workplace	Delivery room	65
		NICU	35
5	Duty shift	Morning	35
		Evening	27
		Night	38

No.: Number, %: Percentage, M: Mean, SD: Standard deviation

Table 2 presents the description of professional variables for nurses and midwives; the findings reveal 7±6 years of experience in general, with 49.1% having 1 – 5 years of experience. The average years of experience in the current unit is 5 ±5, and 61.3% have 1 – 5 years, indicating that nurses and midwives have moderate experience. Regarding participation in training courses about

Kangaroo Mother Care, less than half participated, with most participants attending 1 – 3 training courses. The current workplace shows that 65% of nurses and midwives work in the delivery room and 35% work in the Neonate Intensive Care Unit (NICU). Concerning duty shifts, 38% of nurses and midwives work during the night.

Table 3: Assessment of Nurses and Midwives' Knowledge Items about Implementation of Kangaroo Mother Care (KMC) (N=163)

List	Knowledge	Don't know f (%)	Know f (%)	Mean score	Assess.
1	KMC promotes emotional bonding between the mother and the newborn	12(7.4)	151(92.6)	.93	Good
2	KMC facilitates the stabilization of the newborn's body temperature (thermoregulation)	12(7.4)	151(92.6)	.93	Good
3	The application of skin-to-skin contact (SSC) helps stabilize the newborn's oxygen saturation levels	19(11.7)	144(88.3)	.88	Good
4	KMC promotes and supports exclusive breastfeeding	19(11.7)	144(88.3)	.88	Good
5	KMC enhances the newborn's immunity and reduces the risk of neonatal morbidity	33(20.2)	130(79.8)	.80	Good
6	It is preferable to defer KMC if the newborn is connected to mechanical ventilation	60(36.8)	103(63.2)	.63	Moderate
7	It is preferable to delay KMC for newborns weighing less than 1,000 grams until clinical stability is achieved.	56(34.4)	107(65.6)	.66	Moderate
8	Newborns exhibit signs of comfort and satisfaction during the practice of KMC.	19(11.7)	144(88.3)	.88	Good
9	KMC can be implemented for newborns undergoing phototherapy.	63(38.7)	100(61.3)	.61	Moderate
10	KMC assists in stabilizing the newborn's heart rate and regulating cardiac rhythm.	21(12.9)	142(87.1)	.87	Good
11	KMC can be safely applied to newborns with intravenous (IV) infusion sets.	43(26.4)	120(73.6)	.74	Good
12	The application of KMC regulates the respiratory rate of the neonate.	27(16.2)	136(83.4)	.83	Good
13	It is preferable to avoid KMC for neonates born at less than 28 weeks' gestation until their health status stabilizes.	55(33.7)	108(66.3)	.66	Moderate
14	KMC is safe for medically stable preterm infants	31(19)	132(81)	.81	Good
15	KMC contributes to the reduction of stress and infant crying.	17(10.4)	146(89.6)	.90	Good
16	KMC promotes physiological weight gain in preterm infants.	35(21.5)	128(78.5)	.79	Good

M: Mean, f: Frequency, %: Percentage assessment: Assess, Poor = 0.00 – 0.33, Moderate = 0.34 – 0.67, Good= 0.68 – 1.00

Table 3 presents the assessment of knowledge items among nurse midwives about Kangaroo Mother Care (KMC). The nurses and midwives demonstrated a good level of knowledge regarding KMC's role in reducing infant stress (89.6%), stabilizing heart rate (87.1%), and promoting exclusive breastfeeding (88.3%). However, a knowledge gap was found in aspects

related to complex clinical scenarios, as indicated by a moderate level of knowledge in items related to implementing KMC during phototherapy (61.3%), managing newborns on mechanical ventilation (63.2%), and the care of extremely low-birth-weight infants under 1,000 grams (65.6%).

Table 4: Assessment of Perceived Barriers by Nurses and Midwives Regarding Implementation of Kangaroo Mother Care (KMC) (N=163)

List	Barriers	No f (%)	Yes f (%)	Mean score	Assess.
1	Lack of training or professional knowledge	22(13.5)	141(86.5)	.87	High
2	Shortage of equipment or inadequate space	36(22.1)	127(77.9)	.78	High
3	Absence of health system support for KMC implementation	31(19)	132(81)	.81	High
4	Shortage of healthcare staff in the delivery and Neonatal Intensive Care Units (NICU).	47(28.8)	116(71.2)	.71	High
5	High workload and excessive duties assigned to a single nurse-midwife	32(19.6)	131(80.4)	.80	High
6	Absence of clear hospital policies or clinical guidelines regarding KMC application	40(24.5)	123(75.5)	.75	High
7	Lack of motivation among healthcare staff to practice KMC	36(22.1)	127(77.9)	.78	High
8	Parental refusal or hesitation to participate in KMC.	43(26.4)	120(73.6)	.74	High
9	Difficulty in ensuring maternal privacy during KMC sessions	29(17.8)	134(82.2)	.82	High
10	The belief that conventional incubators are more beneficial than KMC	58(35.6)	105(64.4)	.64	Moderate
11	Fear of accidental ex-tubation or displacement of medical devices during KMC	31(19)	132(81)	.81	High
12	Fear of accidental dislodgement of intravenous (IV) cannulas or infusion sets during KMC.	32(19.6)	131(80.4)	.80	High
13	Time constraints faced by nurses and midwives for KMC implementation	44(27)	119(73)	.73	High
14	Clinical instability of the newborn or excessive crying, limiting KMC feasibility	43(26.4)	120(73.6)	.74	High
15	Traditional newborn care practices (e.g., early bathing or swaddling) hinder KMC	43(26.4)	120(73.6)	.74	High
16	Poor communication and interpersonal dynamics between nurse-midwives and parents	38(23.3)	125(76.7)	.77	High
17	Frequent visitors are limiting the mother's ability to practice KMC.	25(15.3)	138(84.7)	.85	High

M: Mean, f: Frequency, %: Percentage assessment: Assess

Low = 0.00 – 0.33, Moderate = 0.34 – 0.67, High= 0.68 – 1.00

The findings in Table 4 depict that nurses and midwives perceive significant barriers to the implementation of Kangaroo Mother Care (KMC). The biggest barrier is the lack of experience or professional expertise (86.5%, mean = 0.87), which is closely followed by external factors like regular visitor attendance (84.7%) and worries about maternal privacy

(82.2%). Systemic issues also weigh heavily, with a lack of support from health systems (81%) and high workloads (80.4%) cited as having an impact, as well as the fear of dislodgement of medical devices such as IV cannulas or accidental discharge from ventilators being reported (81% and 80.4%, respectively). While most barriers were rated high, the belief that conventional

incubators are superior to KMC was the only item rated as "Moderate" (64.4%), suggesting that while staff values KMC in theory, practical

concerns regarding safety, staffing, and facility infrastructure create a substantial gap between evidence-based knowledge and clinical practice.

Table 5: Overall Assessment of Perceived Barriers regarding Implementation of Kangaroo Mother Care among Nurses and Midwives (N = 163)

Perceived Barriers	f	%	M ± SD	Assessment
Low	1	.6	13.13 ± 3.289	High perceived barriers
Moderate	47	28.8		
High	115	70.6		
<i>Total</i>	<i>163</i>	<i>100</i>		

f: Frequency, %: Percentage

M: Mean for total score, SD: Standard deviation for total score

Low = 0.00 – 5.66, Moderate = 5.67 – 11.33, High = 11.34 – 17.00

Table 5 exhibits that 70.6% of nurses and midwives perceive a “high” level of barriers to the implementation of kangaroo mother care, with a

mean score of 13.13 ± 3.289 . Only 0.6% of them perceived a “low” level of barriers, while 28.8% perceived a “moderate” level of barriers.




Figure (5-1): Levels of Perceived Barriers Regarding Implementation of Kangaroo Mother Care among Nurses and Midwives (N = 163)

This figure shows that 70.6% of nurse midwives perceive high barriers regarding the

Discussion

Socio-Demographic Characteristics

The analysis of the findings shows the overall demographic characteristics of the 163 nurses and midwives who participated in the study. The findings reveal that the highest percentage of nurses and midwives is young. More than half of the nurse-midwives (57.7%) fell within the 20–29 age group, followed by those aged 30–39 years (31.3%).

The current results show a similar pattern to that reported by Ridha and Hanon (2025), who evaluated how an educational program affected nurses' and midwives' understanding of the benefits and barriers of skin-to-skin contact in Kirkuk City hospitals. The study found that over 40% of all nurse-midwives were in the 20–29 age range.

Regarding marital status, nurses and midwives showed that nearly two-thirds were married (65%), while 31.9% were single. Only a small proportion were divorced (1.8%) or widowed (1.2%). The predominance of married participants may be related to the sample's age distribution, as this age group is commonly considered appropriate for marriage in the local community.

Regarding residency, it indicates that the majority (95.7%) of nurses and midwives reside in urban areas, while only 4.3% reside in rural areas. Regarding socioeconomic status, more than half of nurses and midwives reported that their income was sufficient, while a considerable proportion reported it as barely sufficient, and a small proportion reported it as insufficient.

Professional Characteristics

The findings reveal a range of years of experience as a general; 49.1% have 1 – 5 years.

of experience. The years of experience in the current unit are as follows: 61.3% have 1–5 years, indicating that nurse midwives have moderate experience. These findings are in agreement with the study by Kaynat et al. (2024). professional

implementation of kangaroo mother care.

experience (42.8%), indicating a moderately experienced workforce.

Regarding participation in Kangaroo Mother Care training courses, less than half of the participants reported attending such programs, with most of them having participated in only one to three training sessions.

In contrast, a study conducted in Saudi Arabia by Almutairi (2022) reported that a considerable proportion of nurses did not receive adequate training on skin-to-skin contact during their orientation, and many, 42.5%, perceived existing guidelines as unclear or insufficient.

Surveying the current workplace reveals that 65% of nurse midwives work in the delivery room, and 35% work in the Neonatal Intensive Care Unit (NICU).

Regarding duty shift, 38% of nurse midwives work during the night, 35% work during the day (morning), and 27% work during the day (evening).

Knowledge about the Implementation of Kangaroo Mother Care

Nurses and midwives demonstrate a good level of knowledge on the implementation of Kangaroo Mother Care (KMC), according to an assessment of knowledge items among nurses and midwives. Understanding emotional bonding and thermoregulation was found to be highly understood. These results align with the research conducted by Omer (2021).

A cross-sectional descriptive study by Al-Shehri & Binmanee (2021), among NICU nurses in Riyadh, Saudi Arabia, examined the level of the nurses' knowledge about KMC in healthcare facilities. NICU nurses were found to have adequate knowledge. 92.6% of nurse midwives had demonstrated strong knowledge in this regard.

However, a knowledge gap was found in aspects related to complex clinical scenarios, as indicated by a moderate level of knowledge in items related to implementing KMC during phototherapy

(61.3%), managing newborns on mechanical ventilation (63.2%), and the care of extremely low-birth-weight infants under 1,000 grams (65.6%).

This data indicated that there is a need for targeted educational programs regarding the proper execution of KMC for infants who are medically

Conclusion

The study revealed that nurses and midwives face multiple barriers to implementing kangaroo mother care. The main challenges at the individual level include limited clinical competence. Although nurses and midwives understand the importance of KMC, environmental and organizational challenges, such as a lack of privacy for the mother, frequent visitor disruptions, and inadequate institutional support, limit its consistent use in the neonatal unit and a lack of engagement from hospital management.

The study by Cai et al. (2022) found several barriers that hinder the implementation of kangaroo mother care, primarily focusing on the absence of privacy. Furthermore, inadequate leadership, a lack of professional expertise among staff, and insufficient training posed additional challenges for KMC. Systemic issues also weigh heavily, with a lack of support from health systems (81%) and high workloads (80.4%) cited as having an impact.

These findings are congruent with the study by Tumukunde et al. (2024), highlighting significant obstacles such as insufficient space for KMC beds attendance (84.7%) and worries about maternal privacy (82.2%).

Although most barriers were rated high, the perception that conventional incubators are better than KMC scored "Moderate" (64.4%), indicating that, as much as staff believes in KMC in theory, practical factors related to safety, staffing, and facility are such that there is a gap between evidence-based information and its application in clinical care at this NICU.

vulnerable or reliant on technology to achieve completeness.

Perceived Barriers

The biggest barrier is the lack of experience or professional expertise (86.5%), closely followed by external factors such as regular visitor

Recommendation

1. Integrate Kangaroo Mother Care into hospital policies and protocols, and ensure its consistent and sustainable implementation.
2. Implement regular training programs for nurses and midwives to strengthen their knowledge and skills in Kangaroo Mother Care (KMC).
3. Improve the hospital environment by ensuring adequate privacy for mothers during KMC practice. Regulate visiting hours within neonatal units, providing appropriate seating and equipment to support skin-to-skin contact.
4. Encourage continuous monitoring and evaluation of KMC practices to ensure adherence and quality of care.
5. Conduct further research on kangaroo mother care to evaluate its implementation in complex clinical cases such as neonates receiving mechanical ventilation or phototherapy and to identify the organizational barriers to its implementation within healthcare institutions.

References

1. Abadía-Barrero, C. E. (2018). Kangaroo Mother Care in Colombia: A Subaltern Health Innovation against For-Profit Biomedicine. *Medical Anthropology Quarterly*, 32(3), 384–403.
2. Al Mutair, A., Almutairi, W., Aljameez, F., Kay, E., Rabanal, R., Abellar, K., Napaod, A., Hawsawi, A., Saha, C., & Ahmed, G. Y. (2023). Assessment of

- nurses' knowledge, attitude, and implementation of skin-to-skin care within the perinatal setting in Saudi Arabia: Survey study. *Nursing Open*, 10(4), 2165–2171.
3. Alenchery, A. J., Thoppil, J., Britto, C. D., De Onis, J. V., Fernandez, L., & Suman Rao, P. N. (2018). Barriers and enablers to skin-to-skin contact at birth in healthy neonates—A qualitative study. *BMC Pediatrics*, 18(1), 48.
 4. Cai, Q., Chen, D.-Q., Wang, H., Zhang, Y., Yang, R., Xu, W.-L., & Xu, X.-F. (2022). What influences the implementation of kangaroo mother care? An umbrella review. *BMC Pregnancy and Childbirth*, 22(1), 851.
 5. Calibo, A. P., De Leon Mendoza, S., Silvestre, M. A., Murray, J. C. S., Li, Z., Mannava, P., Kitong, J., Quiazon, M. B., & Sobel, H. L. (2021). Scaling up kangaroo mother care in the Philippines using policy, regulatory, and systems reform to drive changes in birth practices. *BMJ Global Health*, 6(8), e006492.
 6. Chamhene, N., & Moshi, F. V. (2023). Level of Knowledge on Kangaroo Mother Care and Its Associated Factors Among Postnatal Mothers With Preterm Babies in Central Tanzania: Hospital-Based Cross-Sectional Study. *SAGE Open Nursing*, 9, 23779608231167813.
 7. Darghiyan, Z., Ghasemi, F., Karami, K., Valizadeh, F., & Mohammadi, R. (2023). A comparative study of the effects of kangaroo care by mothers and maternal grandmothers on the vital signs of hospitalized preterm newborns: A randomized controlled clinical trial study. *Trials*, 24(1), 275
 8. Kaynat, C., Saoul, A., Shehzad, H., Ullah, K., Tasneem, S., & Jabeen, R. (2024). Assessment of knowledge regarding kangaroo mother care among nurses of a general hospital in Lahore. *Biological and Clinical Sciences Research Journal*,
 9. Liu, T., Xu, Y., Gong, Y., Zheng, J., & Chen, Z. (2024). The global burden of disease attributable to preterm birth and low birth weight in 204 countries and territories from 1990 to 2019: An analysis of the Global Burden of Disease Study. *Journal of Global Health*, 14, 04109.
 10. Mala, S. A., & Abdul-Sahib, S. H. (2022). The Impact of Nurse-Related Barriers on Kangaroo Mother Care Implementation in Kurdistan of Iraq. *Kufa Journal for Nursing Sciences*, 12(2).
 11. Maniago, J. D., Almazan, J. U., & Albougami, A. S. (2020). Nurses' Kangaroo Mother Care practice implementation and future challenges: An integrative review. *Scandinavian Journal of Caring Sciences*, 34(2), 293–304.
 12. Ridha, N. R., & Hanon, R. H. (2025). Effect of an educational program on knowledge and perceived barriers toward kangaroo care among nurses and midwives in Kirkuk hospitals. *Current Clinical and Medical Education*, 3(7), 29–40
 13. Omer, N. M. (2021). Assessment Of Nurses' and Midwives' Knowledge, Beliefs, and Barriers Regarding Kangaroo Care in Erbil. Master's Thesis in the Nursing Program of the Near East University Institute ...
 14. Sales, I. M. M., Santos, J. D. M., Rocha, S. S. D., Gouveia, M. T. D. O., & Carvalho, N. A. R. D. (2018). Contributions of the nursing team in the second stage of the Kangaroo-Mother Care Method: Implications for hospital discharge of the newborn. *Escola Anna Nery*, 22(4).
 15. Tumukunde, V. S., Katongole, J., Namukwaya, S., Medvedev, M. M., Nyirenda, M., Tann, C. J., Seeley, J., & Lawn, J. E. (2024). Kangaroo mother care before clinical stabilization: Implementation barriers and facilitators reported by caregivers and healthcare providers in Uganda. *PLOS Global Public Health*, 4(7), e0002856.

16. UNICEF. (2022). Neonatal mortality rate in Iraq. UNICEF Data. <https://data.unicef.org>
17. Gebeyehu, N. A., Gelaw, K. A., Azeze, G. A., Admass, B. A., Lake, E. A., & Adela, G. A. (2022). Knowledge, attitude and practice towards kangaroo mother care among postnatal women in Ethiopia: Systematic review and meta-analysis. PLOS ONE, 17(5), e0265411.
18. Chan, G., Bergelson, I., Smith, E. R., Skotnes, T., & Wall, S. (2017). Barriers and enablers of kangaroo mother care implementation from a health systems perspective: A systematic review. Health Policy and Planning, 32(10), 1466–1475.
19. Al-Shehri, H., & Binmanee, A. (2021). Kangaroo mother care practice, knowledge, and perception among NICU nurses in Riyadh, Saudi Arabia. *International Journal of Pediatrics and Adolescent Medicine*, 8(1), 29–34.